

Genetic Identification of Necrophagous Insect Species (*Diptera*) of Forensic Importance Sampled from Swine Carcasses in Mato Grosso, Midwestern Brazil

Elisângela Santana de Oliveira Dantas^{3,4}, Diniz Pereira Leite Junior^{1,4*}, Junio Damasceno de Souza⁵, Reginaldo Rossi do Carmo², Flavia Galindo Silvestre Silva² and Patricia Pasquali Parisi Maltempi³

¹UFMT – Federal University of Mato Grosso - Av. Fernando Correa da Costa, Brazil

²POLITEC – Official and Technical Expertise Identification, Government of Mato Grosso - Av. Gonçalo Antunes de Barros, Carumbe, Brazil

³UNESP – State University of Sao Paulo, Institute of Biosciences, Rio Claro Campus, Technical Section Post Graduate, Brazil

⁴UNIVAG – University Center of Varzea Grande, Av. Dom Orlando Chaves, Brazil

⁵UniBH - Centro Universitario de Belo Horizonte, PUC Minas, Rua Diamantina, Brazil

*Corresponding author: Junior DPL, Federal University of Mato Grosso, Faculty of Medicine, Research Laboratory, Av. Fernando Correa da Costa, 2367, Boa Esperanca – Cuiaba, MT 78060-900, Brazil, Tel: +55(65)3615-8809; E-mail: djbiologico@gmail.com

Received date: Feb 21, 2016; Accepted date: May 09, 2016; Published date: May 12, 2016

Copyright: © 2016 Dantas ESO, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Forensic entomology is the science that uses insects found on bodies to set the Post Mortem Interval (PMI) among other investigations. The present work used DNA barcode based on the use of a standardized fragment of Cytochrome C Oxidase subunit I (COI). We undertook analysis of the taxa associated with carcasses, representing 15 dipterans species. Carcasses of pigs during the decomposition process were used, using a cutoff value of 3% for intra- and interspecific differentiation proves efficient for the dipteran of forensic interest. Species identification is more accurate in well studied and reviewed groups. It was observed that specific taxonomic groups were well separated. In the larger group a molecular study, including other mitochondrial genes and nuclear genes is necessary. The greater differentiation in *Lucilia eximia* probably requires review studies of the group.

Keywords: Forensic science; Forensic entomology; Barcode; DNA; *Diptera*; Mato grosso

Introduction

Necrophagous insects are ubiquitous insects occurring in anthropogenic ecosystems. Among them we can highlight the flies which have great medical importance due to their participation in carrion decomposition, facultative parasitism of vertebrate tissues and mechanical transmission of various pathogenic microorganisms [1-3]. In recent decades, there has although there are insect knowledge of the use records for forensic application, historically, the early works were influenced by studies by Jean Pierre Megnin in the second half of the nineteenth century [4]. Currently several publications on the subject has been discussed by several authors [5-8]. But in the center west of Brazil, the state of Mato Grosso; this issue is rarely addressed. This research is the initial registration of this activity.

After a lengthy hiatus of over 50 years new studies were produced propelled by the rise of new research groups in forensic entomology since 1991 [9]. Currently in Brazil, more than 20 scientists and dozens of forensic experts with field training are carrying out research projects related to forensic entomology spread along all Brazilian states [10]. Ironically, Brazilian cities suffering from the highest rates of violence have been neglected in field inventories? Of forensically important species [11,12]. Partly due to convoluted mechanisms of Brazilian legal system and lack of investment in intelligence procedures for the police modus operandi [10].

A variety of necrophagous insect species occurs on or around cadavers and carcasses depending on their preference for a given stage of decomposition [13]. Systematized and pioneer studies of cadaver

entomofauna were published in Mégnin's book [14]; the author listed the chronological sequence of body decomposition and insect colonization. The colonization sequence was described as eight waves of invasion whose we were able to recognize the orders: *Diptera*, *Coleoptera*, *Lepidoptera* and *Acarina*.

According to Oliveira-Costa [15], the entomological succession follows an initially predictable pattern since every phase of decomposition processing the stratum is colonized by specific groups which in turn, attract different stages making it possible to associate specific species to specific phases of the decomposition process.

Insects can be useful for crime investigation because they provide an estimation of the time elapsed after death known as Post Mortem Interval (PMI). PMI is based on the time which an insect takes to develop from an egg to the stage in which it is found on the body. After identification of the cadaveric fauna several clues about the crime can be inferred by linking these bugs or fragments found on victim such as where the crime took place and the suspects of the crime [16-18]. Due to their keen sense of smell dipterans rapidly locate dead bodies by their scent then colonizes this resourceful habitat as a reproduction site for oviposition also for nutritional source [16,19,20].

There are studies conducted in other countries that combine a large database on the succession pattern of insects on bodies. However, due to differences in climate conditions along with huge territorial extension, these data cannot be safely used in our forensic examinations [21].

Therefore, the present study has the objective of describing *Diptera* entomological species succession in different carcasse decomposition phases on climate conditions of Cuiaba/MT. We used DNA barcode to

assemble an easy access database to be utilized by forensic experts in criminal investigations.

Materials and Methods

Sample collection

The study was conducted in an area inside POLITEC. The public institution responsible for official forensic examinations in the state of Mato Grosso, Brazil (GPS: L 06-01-88-4/N 82-77-22-8). As substrate, we used swine carcasses *Sus scrofa* (*Suidae*) inside a metallic cage (60 × 90 × 45 cm high) until their complete decomposition.

Experimental models of carcasses used in the research were large white domestic pigs with a body weight of about 15-20 kg. Carcasses of pigs were acquired through traders and merchants who slaughtered pigs in farms near the city of Cuiaba. The animals were slaughtered at 06:00 am and brought to the worksite two hours late where the decomposition process and consequently the collection of specimens *Diptera* later occur. The requirements on ethics permits were not required at the time of the experiment and the ethical committees were not available for any evaluations accordingly; as the case of a legal and common practice in the local trade fairs in the state of Mato Grosso.

The climate is characteristic of the semiarid region, with a mean annual rainfall of 1469.4 mm and average annual temperature of 25°C to 40°C. Despite this inequality, the region is well supplied with rain and seasonality is typically tropical, with maximal temperatures in summer and minimal in winter.

Four experiments were conducted over the seasons. One experiment for each season of the year between January 2013 and December 2013. The region has a hot sub-humid tropical climate with the summer high temperature and the winter minimum temperature around 17°C. The winters are excessively dry because rainfall is very rare. Spring and fall are unstable. The study area are surrounded by open arboreal vegetation.

The phases of decomposition were observed until complete skeletonization. Where decomposition phases being defined as fresh, bloat, active decay, advanced decay and dry remains phase. The carcasses were monitored every day at 08:00 a.m. and 10:00 a.m. until complete decomposition of the individual. For entomofauna collection, larvae was collected with a tablespoon every three days in three different regions of the animal choosing only well-developed bigger larvae.

For identification part of collected specimens was transported alive to the forensic entomological lab at POLITEC where larvae completed their life cycles necessary for identification through morphology. A second part of larvae were fed for three days in plastic containers containing sawdust. After that they were washed in water and stored in 96% ethanol in a -20°C freezer. This material was stored and then used in following molecular processes.

DNA extraction and molecular analysis

The DNA was extracted only from the immature stages. The extractions were carried out from a whole individual, as follows: For the Mix we used 60 µl of EDTA (C=0.5M; pH 8.0) and 250 µl of Nucleus Lysis Solution for each sample. The Mix was chilled on ice becoming cloudy and gelatinous. We distributed the Mix in big eppendorf tubes and macerated the individual larvae. We added 10µl of Proteinase K (20 mg/ml) to the solution and agitated it. The tubes

were left in a water bath at 65°C for 2 hours. It was optional to add 1.5 µl of RNase to the solution, mixing the sample by inverting the tube 2-5 times. We incubated the samples at 37°C for 30 minutes. After, we let the samples cool to room temperature for 5 minutes. 100µl of Protein Precipitation Solution was added (Promega) and vortexed for 20 seconds. The samples were chilled on ice for 5 minutes. We centrifuged the samples for 10 minutes at 13,000-16,000×g. The protein will form a white pellet. We transferred the DNA containing supernatant to a large eppendorf already containing 300 µl of isopropanol (room temperature). We mixed the solution inverting the tubes and centrifuged it for 2 minutes at 13,000-16,000×g. DNA will form a white pellet. We carefully removed the supernatant. We added 300 µl of 70% Ethanol and carefully inverted the tube several times to wash the DNA. We centrifuged the sample again for 2 minutes at 13,000-16,000×g, carefully removing the supernatant after that. We left the samples in a water bath (40°C) for 30 minutes in order for the alcohol to evaporate. We added 60-100 µl of autoclaved Milli-Q H₂O. The samples were, then, incubated overnight at 4°C (or room temperature). We briefly and carefully vortexed the samples and then store them in a freezer.

DNA from whole specimens stored in 96% ethanol was extracted. The samples were macerated in lysis solution containing SDS/ Proteinase K (480 µl of 50 mM Tris, 50 mM EDTA, pH 8.0; 20 µl of 20 mg/mL proteinase K) and incubated for 3 hours in a water bath at 55°C. Afterwards, we added 500 µl of EZDNA KIT (Biosystems[®]) followed by a new incubation at room temperature for 12 hours. The samples were centrifuged and the supernatant was washed 3 times in 95-100% ethanol for DNA precipitation. After the alcohol evaporated, DNA was resuspended in 70 µl of TE (10 mM Tris, 50 mM EDTA, ultrapure H₂O), quantified in a spectrophotometer and stored in a freezer.

Amplification

The mitochondrial DNA region analyzed was the final portion of the cytochrome c oxidase subunit I (COI), corresponding to the Barcode DNA region [22]. This segment was amplified through the initiators set LCO1490-L (18 µl) and HCO2198-L (18 µl) [23]. Amplification was performed using: 40 ng of DNA, 25 mM of dNTP, 20 mM of each initiator, 2.5 U of Taq and 3-4 mM of MgCl₂ (Biosystems). PCR temperature cycles consisted in an initial denaturation step at 94°C for two minutes, followed by 30 denaturation cycles at 94°C for 30 seconds, annealing at 46°C for 30 seconds and elongation at 72°C for 2 minutes. The last cycle was followed by a 5 minute incubation at 72°C for a final elongation [23].

Sequencing

Amplicons were visualized in a 1.5% agarose gel electrophoresis, stained with ethidium bromide. Then, purified according to protocol based on PolyEthylene Glycol (PEG) and stored in a freezer. Sequencing reaction was carried out with 40 ng of DNA; 1.6 µM of the same PCR initiator and specific sequencing kit.

Amplification was conducted in a thermocycler in which cycles consisted of initial denaturation at 96°C for 1 minute, 35 denaturation cycles at 96°C for 10 seconds, annealing at 50°C for 5 seconds and elongation at 60°C for 4 minutes [23]. Sequencing was done at Politec-MT forensic laboratory.

Data entry in the database

Once we obtained the genetic sequence of the questioned sample, analysis and editing are necessary for submitting the sequence to the database for similarity search. Gross data from the sequencer are called base calling.

Allignment

Consensus sequences were aligned in Clustal W software [8] using default parameters then manually checking and editing them. Blast's major utility in this study was to compare between nucleotides sequences provided by the user (query) with other sequences inside the database. Sequence comparison was analyzed through pair by pair alignment where every nucleotide pair receives a score determined by a score matrix in which values are empirically determined. When there is no gap in a sequence it is represented in a graph by a continuous diagonal line. Otherwise that line would be interrupted one or more times [24].

Result

Of the 220 individuals initially selected only 50% had their DNA amplified by the COI region. This because their supporting amounts, result in a 50% success rate amplification. Because of the low value of the core samples at the time of recognition of the species; generating ambiguous samples. The ones that were successfully amplified 13 species could be identified through 613 bp sequences.

This systematic approach provides the initial infrastructure for the next generation of biodiversity assessment and environmental monitoring adults of necrophagous blowflies. It can lead to more effective understanding, appreciation, and management of these complex ecosystems. A geographical distribution of the species is shown in Table 1.

Super-family /Family	Subfamily	Tribe	Genus	Species	Geographical distribution
Muscoide Muscidae	Muscinae	Muscini	<i>Musca</i>	House fly <i>M. domestica</i> Linnaeus, 1758 [29]	Cosmopolitan
	Azeliinae	Azeliini	<i>Ophyra</i>	Dung fly <i>O. albuquerquei</i> (Loper, 1985)	Brazil, Argentina
	Azellinae	Azeliini		<i>O. aenescens</i> (Wiedemann, 1830)	USA, Nicaragua, Bahamas, Bermuda, Greater and Lesser Antilles, French Polinesia, Hawaii, Neotropical Region
Oestroidea Calliphoridae	Toxotarsinae	Sarconesiini	<i>Sarconesia</i>	Blow fly <i>S. chlorogaster</i> (Wiedemann, 1831)	Peru, Brazil, Paraguay, Uruguay, Argentina, Chile, Juan Fernandez Islands and Easter Islands
	Chrysomyinae	Chrysomyini	<i>Chrysomya</i>	<i>C. albiceps</i> (Wiedemann, 1819)	Africa, Mediterranean, North American and South American and Brazil
				<i>C. megacephala</i> (Fabricius, 1794)	Africa, Australasia, Palearctic and Oriental regions until Southern Africa and Afrotropical islands
				<i>C. putoria</i> (Wiedemann, 1818)	Argentina, Bolivia, Brazil, Colombia, Panama, Paraguay, Peru
	Calliphorinae	Luciliini	<i>Lucilia = Phaenicia*</i>	Green-bottle fly <i>L. sericata</i> (Meigen, 1826)	Southern Canada, Argentina, Bermuda, Brazil, Chile, Colombia, Peru, Venezuela
				<i>L. cluvia</i> (Walker, 1849)	Southern USA, Cuba, Puerto Rico, Bahamas, Argentina, Colombia, Guatemala, Honduras, Martinique, Mexico, Nicaragua
				<i>L. porphyrina</i> (Walker, 1856)	Sri-Lanka, China, India, Malasya, Nepal and Philipines

				<i>L. cuprina</i> (Wiedemann, 1830)	Tropical Regions, southern USA, Argentina, Brazil, Colombia, Cuba, Haiti, Jamaica, Peru, Uruguay, Venezuela
				<i>L. eximia</i> (Wiedeman, 1819)	Southern USA, Mexico, Peru, Venezuela, Brazil, Argentina, Bahamas, Nicaragua, Chile, Colombia
				<i>L. mexicana</i> (Macquart, 1843)	Southern USA, Mexico, Guatemala, southern Brazil
				<i>L. coeruleiviridis</i> (Macquart, 1855)	U.S.A., Maryland, Baltimore, Neotropical: Cuba, Guatemala
Oestroidea Sarcophagidae	Sarcophaginae	Sarcophagini	Peckia (<i>Pattonella</i>)	Flesh fly <i>P. intermutans</i> (Walker, 1861)	concentrated in regions of tropical and warm temperate climate and in Neotropical region
This information and classification is modeled on the basis of previous reports by Kosmann et al. [51]; ITIS Report [50]*; Whithworth [25] and Rognes [52], respectively.					

Table 1: Necrophage insects collected in swine (*Sus scrofa* L.) carcasses in Cuiabá/MT region.

Species		1	2	3	4	5	6	7	8	9	10	11	12	13	
		0.01	0.009	0.009	0.009	0.01	0.011	0.009	0.009	0.008	0.014	0.014	0.013	0.012	
<i>L. cluvia</i>	0.074		0.007	0.011	0.011	0.008	0.009	0.012	0.011	0.011	0.014	0.015	0.014	0.013	
<i>L. coeruleiviridis</i>	0.056	0.03		0.009	0.009	0.007	0.008	0.011	0.009	0.009	0.013	0.013	0.013	0.013	
<i>L. cuprina</i>	0.05	0.08	0.062		0.001	0.01	0.011	0.01	0.003	0.008	0.015	0.013	0.012	0.011	
<i>L. cuprinahawaii</i>	0.05	0.08	0.061	0.001		0.01	0.011	0.01	0.003	0.008	0.015	0.013	0.012	0.011	
<i>L. eximia</i>	0.084	0.06	0.048	0.082	0.081		0.005	0.012	0.01	0.01	0.014	0.013	0.013	0.012	
<i>L. mexicana</i>	0.074	0.05	0.044	0.073	0.072	0.032		0.012	0.011	0.011	0.015	0.014	0.014	0.013	Stand ard deviat ion
<i>L. porphyrina</i>	0.055	0.09	0.079	0.071	0.071	0.1	0.096		0.011	0.01	0.015	0.015	0.014	0.014	
<i>L. sericata</i>	0.049	0.08	0.058	0.007	0.006	0.082	0.072	0.074		0.008	0.015	0.013	0.012	0.011	
<i>L. thatuna</i>	0.044	0.08	0.054	0.038	0.038	0.077	0.07	0.069	0.039		0.013	0.013	0.013	0.011	
<i>M. domestica</i>	0.105	0.12	0.103	0.123	0.123	0.135	0.136	0.13	0.12	0.107		0.013	0.014	0.013	
<i>O. aenescens</i>	0.113	0.13	0.109	0.106	0.107	0.133	0.127	0.128	0.103	0.117	0.103		0.01	0.014	
<i>O. albuquerquei</i>	0.107	0.13	0.103	0.093	0.095	0.13	0.121	0.124	0.09	0.098	0.124	0.063		0.012	
<i>S. chlorogaster</i>	0.086	0.11	0.097	0.081	0.081	0.116	0.109	0.116	0.077	0.073	0.106	0.123	0.101		
Genetic difference															

Table 2: Genetic distance of the analyzed species.

The present study is the first to report the analysis of the obtained results made it possible to verify the succession pattern of these species in the decomposition process of pigs (*Sus scrofa*) (Table 2).

Members of the family *Calliphoridae* were the most commonly isolated. This genus includes numerous *saprophagous* species formerly *Phaenicia* [25]. A common visitor to carrion, feces and garbage plays an important role in forensics, medical science and veterinary medicine.

In the fresh phase, *Calliphoridae* was dominant, especially *Lucilia eximia* (Wiedemann, 1819) followed by *Chrysomya megacephala* (Fabricius, 1794) and *Musca domestica*. In the bloat phase, we observed *L. eximia*, *C. albiceps* (Wiedemann, 1819), *C. putoria* (Wiedemann, 1818), *Peckia (Pattonella) intermutans* (Walker, 1861), *Ophyra aenescens* (Wiedemann, 1830) and *O. albuquerquei* (Lopes, 1985). During active decay phase we observed immatures of: *C. megacephala*, *C. albiceps*, *C. putoria* (Wiedemann, 1818), *P.*

intermutans, *M. domestica*, *L. cuprina* (Wiedemann, 1830), *L. sericata* (Meigen, 1826), *L. cluvia* (Walker, 1849), *L. coeruleiviridis* (Macquart, 1855), *L. mexicana* (Macquart, 1843) and *L. porphyrina* (Walker, 1856). In advanced decay phase, it was possible to observe *C. albiceps*, *C. megacephala*, *Sarconesia chlorogaster* (Wiedemann, 1830) and *M. domestica*. During the dry remains phase, we could observe *M. domestica*, *C. megacephala* and *C. albiceps*. Other insects from other orders, like: *Coleoptera*, *Hemiptera* and *Dermaptera* were predominant during this period.

With respect to the succession patterns observed in the different seasons of the year we could evaluate that they are extremely similar varying only with regard to the time duration of the fresh and dry remains phases. During the rainy season (November to March) complete decomposition occurred in twelve days. On the other hand, during dry season (April to September) this process was accelerated with total decomposition reaching an average of nine days.

With sequence alignment obtained from COI we obtained a 613 bp sequence matrix. from which genetic distance analysis was carried out using the Tamura-Nei method [26]. Purine and pyrimidine frequency composition analysis for COI segments is summed up in Table 3.

SPECIES	T	C	A	G	Total
<i>L. caeser</i>	37.7	16	30.2	16.6	613
<i>L. coeruleiviridis</i>	37.4	17	29.2	16.5	613
<i>L. cuprina</i>	37.5	16	30.5	16.5	613
<i>L. cuprina hawaii</i>	37.5	16	30.3	16.5	613
<i>L. eximia</i>	37.2	17	28.7	17.1	613
<i>L. mexicana</i>	36.1	18	29.2	16.7	612
<i>L. porphyrina</i>	35.6	18	30.3	16.5	613
<i>L. sericata</i>	37.8	15	30.5	16.3	613
<i>L. thatuna</i>	37.8	15	30.3	16.5	613
<i>M. domestica</i>	38.2	15	29.5	17	613
<i>O. aenescens</i>	36.1	18	29.2	16.5	613
<i>O. albuquerquei</i>	38.7	15	30.3	16.2	613
<i>S. chlorogaster</i>	39.3	14	29.7	16.6	613
Average	38	16	30	16.5	613

Table 3: Purine and pyrimidine frequencies in COI segments.

Discussion

In study area the average annual rainfall is 1.300 mm to 1.450 mm, with maximum intensity in January, February and March. The mean maximum temperature reaches 25°C to 35°C. But the absolute maximum can reach 40C in the warmer months; on rainy days. These meteorological data confirm that the climate of Cuiaba is Cwa. according to the climatic classification of Köppen-Geiger created in 1928 [27]. A Cwa climate presents subtropical characteristics, with a dry winter and a rainy summer.

The Order *Diptera* is divided into two suborders: *Nematocera* which includes mosquitoes along other flies with long antennae and

Brachycera covering the robust and compact flies with short antennas. Among the *Brachycera*, scavenger species of the families *Calliphoridae*, *Sarcophagidae*, *Muscidae*, *Fanniidae* and *Stratiomyidae* are among the representatives of most forensic relevance [7,28,29].

DNA barcoding has become a fairly useable method of choice for rapid species identification in the last decade. Due to the existence of a taxonomic crisis (relative to taxonomic identification of immature specimens that can result in wrong identifications) of the bar code to identify eukaryotic DNA, based on use of a standardized cytochrome fragment (COI); this method has been the subject of much debate [30-33]. The DNA barcoding; It is still the subject of numerous criticisms of its applicability. As opposed to the use of morphology and identification of species and other taxa. Regarding this regard. the identification of forensic interest dipteran. in this work was to record the relevant samples *Calliphoridae* and *Muscidae*, becoming 13 identified species.

Among these proposals, DNA barcoding has been particularly successful in the identification and delineation of new species of various groups and has received greater acceptance because it is simple and affordable [34-38]. Species identification of the adults of necrophagous blowflies is usually not difficult. and numerous keys are currently available for potential users [25,39-42]. However, some characters must nonetheless be treated with caution.

The approach for intra and interspecific differentiation using a 3% cutoff value has shown to be valid for forensically important *diptera*: *Ophyra albuquerquei* and *O. aenescens* as well as *Sarconesia chlorogaster*, *Lucilia thatuna* (Shannon, 1926), *L. cluvia*, *L. porphyrina* and *L. Caesar* (Linnaeus, 1758) [29] were separated from the rest. Gender differences were higher than 3% in all cases. But in some cases, mitochondrial DNA sequence monophilia analyzed here was not proved. It was not possible to separate *L. cuprina* and *L. sericata* that behaved as if they belonged to the same species, as well as *L. coeruleiviridis* and *L. mexicana* that were grouped with different haplotypes comparing to *L. eximia*.

Species identification is more accurate in well studied and revised groups. Interspecific differences varied between 13.6% for *L. mexicana* and *M. domestica* and 0.1% for *L. cuprina* and *L. cuprina hawaii*. Regarding intraspecific differences. *L. eximia* has more than 3% differentiation. Despite the COI gene alone is not being enough for a deep phylogenetic study. It gave support for the formation of important clades. We observed that the specific taxonomic groups were generally well separated. In the larger group where we find: *L. eximia*, *L. Mexicana*, *L. cluvia* and *L. coeruleiviridis* a molecular phylogenetic study including more mitochondrial and nuclear genes might be necessary. Still, *L. Mexicana*, *L. cluvia* and *L. coeruleiviridis* are well separated and have a high support index. The high differentiation in *L. eximia* would probably require a group review in future studies.

Diptera constitute along with *Coleoptera* the majority of the necrophage insects representing the largest portion of cadaveric fauna found on carcasses during the decomposition process [43-45] and that was also observed in this study. For this reason, members of *Sarcophagidae*, *Calliphoridae* and *Muscidae* family have been commonly cited in forensic entomology studies [46,47] as observed in this study as well. Adults of the *Calliphoridae* family are the first to colonize the carcasses due to their agility in laying their eggs soon after the death of the animal. In accordance with this study, members of this family have been frequently observed by other authors [45,46,48,49].

The presence of *M. domestica* and *C. megacephala* during the fresh phase (first 18 hours after death) was also reported by Carvalho and researchers [50] that worked with the succession pattern of insects on dead pigs in the state of São Paulo. Chin and collaborators [51] in their studies are found *Calliphoridae* and *Muscidae* visiting carcasses, finding *C. megacephala* as the dominant species. In our study, we found calliphorid fly (*Sarconesia spp.*, *Chrysomya spp.*, and *Lucilia spp.*) and muscid fly (*Ophyra spp.* and *M. domestica*) visiting the floating carcass; with the gender *Lucilia* as the dominant species. The presence of *L. eximia* during this phase corroborates the data obtained by Souza and researchers [48] as well as its presence in all seasons. The results of the fly species found in this study, approach the species identified in the study by Oliveira-Costa et al. [7] in an experiment in the city of Rio de Janeiro, Brazil. More recently, Salem et al. [52] report on their studies in Egypt, the results of *Chrysomyinae* specimens belonging to three forensically important.

The significant presence of members of the *Calliphoridae* family during the bloat phase corroborates the data from Gomes et al. [53] that already proposed their use as an indirect “biological clock” capable of indicating the amount of time elapsed since death to the Discovery of the body. Members of the *Sarcophagidae* family have been observed by Barros et al. [54] during the period of active and advanced decay phases corroborating the data found in the literature. However, different from what Carvalho and Linhares [46] and Barros et al. [54] observed. The abundance of members from *Sarcophagidae* family was small when compared with members of the *Calliphoridae* family. In 2015, Barbosa and colleagues [55] showed the geographical distribution and diversity of the expansion of the flesh flies distributed by the northeast region of Brazil.

In contrast, in northern Brazil amazonense authors [8] studied the composition and temporal distribution one of the most forensic importance insect families. More currently Akbarzadeh and colleagues [56] identified species of *Calliphoridae* of forensic importance. Showing that in recent decades there has been growing interest in research into the biology and ecology of insects with potential forensic applications [13,56]. Therefore, it must be considered that the active decay phase for the decomposition process in Cuiaba region, starts in the third day after death, for the rainy season, and second day after death for the dry season.

The faster decomposition during the dry season when compared to the rainy season has already been reported by Carvalho & Linhares [45] and Oliveira-da-Silva et al. [57-60] that justify it by saying it happens due to the high temperature, as well as the air's relative humidity. However, total decomposition period is different from that observed by the authors mentioned above, who had periods of 24 days for the rainy season and 11 days for the dry season.

In this study, the use of democratic barcode nurtures the easy access of species identification even when dealing with insect fragments. This work is relevant due to taxonomic identification of insects of forensic interest providing additional information needed to further comprehensive study of these scavengers organisms.

Even so when secure opportunities present themselves safe. we encourage future studies and other researchers to continue documenting species of necrophagous dipterous taking place in this part of the world to help facilitate the various applications of these important insects, in view of the species diversity and overlapping habitat of such flies in Mato Grosso state.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

1. Hall M, Wall R (1995) Myiasis of humans and domestic animals. *Adv Parasitol* 35: 257-334.
2. Greenberg B (1973) Flies and diseases. Biology and disease transmission. Vol. 2. Princeton: Princeton University.
3. Norris KR (1965) The bionomics of blowflies. *Annual Revista Entomologia* 10: 47-68.
4. Carrera M, Insects of medical importance (1991) Publisher of the Federal University of Parana p: 228.
5. Iancu L, Sahlean T, Purcarea C (2015) Dynamics of Necrophagous Insect and Tissue Bacteria for Postmortem Interval Estimation During the Warm Season in Romania. *J Medical Entomology* 53: 54-66.
6. Shin SE, Lee HJ, Park JH, Ko KS, Kim YH, et al. (2015) The First Survey of Forensically Important Entomofauna Collected from Medicolegal Autopsies in South Korea. *BioMed Research International*.
7. Oliveira-Costa J, Lamego CMD, Couri MS, Mello-Patiu CA (2014) Differential Diptera succession patterns onto partially burned and unburned pig carrion in southeastern Brazil. *Brazilian Journal of Biologica* 74: 870-876.
8. Ururahy-Rodrigues A, Rafael JA, Pujol-Luz JR (2013) Temporal Distribution of Blowflies of Forensic Importance (Diptera: Calliphoridae). In *Man-Size Domestic Pigs Carcasses, in the Forest Reserve Adolpho Ducke, Manaus, Amazonas, Brazil*. *Entomobrasilis* 6: 9-22.
9. Salviano RJB, Mello RP, Santos RF, Beck LCNH, Ferreira A (1994) Dipteran caliptrados occurrence in human body. XX Brazilian Congress of Zoology. Rio de Janeiro: RJ.
10. Pujol-Luz JR, Arantes LC, Constantino R (2008) One hundred years of forensic entomology in Brazil (1998-2008). *Journal of Entomology* 52: 485-492.
11. Waiselfisz JJ, Mapa da Violência (2015) Deaths by firearms. Brasília / DF - UNESCO p: 112.
12. Vasconcelos SD, Araújo MCS (2012) Necrophagous species of Diptera and Coleoptera in northeastern Brazil: state of the art and challenges for the Forensic Entomologist. *Revista Brasileira de Entomologia* 56: 7-14.
13. Tomberlin JK, Mohr R, Benbow ME, Tarone AM, Vanlaerhoven S (2011) A roadmap for bridging basic and applied research in forensic entomology. *Annual Review Entomology* 56: 401-421.
14. Mégnin P (1894) *Wildlife Cadaveres*. Paris: Scientific Encyclopedia reminders.
15. Oliveira-Costa J (2011) *Forensic Entomology: when insects are traces*.
16. Miranda G Jacques G, Almeida MP, Silva MSB (2006) Insect samples collection for forensic purposes. Brasília. Justice ministry pp: 4-11.
17. Goff ML, Flynn M (1991) Determination of postmortem interval by arthropod succession: A case study from the Hawaiian islands. *JFSCA* 36: 607-614.
18. Galloway A, Birkby WH, Jones A, Henry TE, Parks B (1989) Decay rates of human remains in arid environment. *JFSCA* 345: 607-616.
19. Greenberg B (1985) *Forensic Entomology: Case Studies*. *ESA* 31: 25-28.
20. Jirón LF, Vargas LG, Vargas-Alvarado E (1983) Four muscoid flies (*Sarcophagidae* and *Muscidae*) associated with human cadavers in Costa Rica. *Brenesia* 21: 3-5.
21. Mann RW, Bass WM, Meadows L (1990) Time since death and decomposition of the human body: variables and observations in case and experimental field studies. *JFSCA* 35: 103-111.
22. Hebert PDN, Cyminska A, Ball SL, Waard JR (2003) Biological identifications through DNA barcodes. *Proceedings of Royal Society of London. Biological Sciences* 270: S569-S599.

23. Nelson LA, Wallman JF, Downton M (2007) Using COI barcodes to identify forensically and medically important blowflies. *Medical and Veterinary Entomology* 21: 44-52.
24. Cruz LM (2006) Blast: theory and practice presented in: I Day of Paraná Bioinformatics– JOBIP, Curitiba.
25. Whitworth T (2005) Keys to the genera and species of blow flies (Diptera: Calliphoridae) of America north of Mexico. *Proceeding of the Entomological Society Washington* 108: 689-725.
26. Tamura K, Nei M (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* 10: 512-526.
27. Rolim G, Camargo MBP, Lania DG, Moraes JFL (2007) Climate classification Köppen and Thornthwaite and its applicability in determining agro-climatic zones in the State of São Paulo. *Bragantia Campinas* 66: 711-720.
28. Gennard D (2012) *Forensic entomology: an introduction*. Oxford: John Wiley and Sons Ltd 2: 272.
29. Fontoura P, Oliveira-Costa J, Celino TB, Costa CES, Rosa DO (2009) Checklist of Diptera of immature associated with pig carcasses *Sus scrofa* (Linnaeus 1758) decomposing in Rio de Janeiro. *Congress Brazil Ecology. St. Lawrence. Proceedings of the IX Congress of Brazil Ecology. St. Lawrence*.
30. Moritz C, Cicero C (2004) DNA barcoding: promise and pitfalls. *Plos Biol* 2: e354.
31. Prendin L (2005) Comment on "Identifying spiders through DNA barcodes". *Canadian Journal of Zoology* 83: 481-491.
32. Meier R, Shiyang K, Vaidya G, Ng PK (2006) DNA Barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. *Systematic Biology* 55: 715-728.
33. Pires AC, Marinoni L (2010) DNA barcoding and traditional taxonomy unified through Integrative Taxonomy: a view that challenges the debate questioning both methodologies. *Biota Neotropica* 10: 339-346.
34. Padial JM, De La Riva I (2007) Integrative taxonomists should use and produce DNA barcodes. *Zootaxa* 1586: 67-68.
35. Smith MA, Rodrigues JJ, Whitfield JB, Deans AR, Janzen DH, et al. (2008) Extreme diversity of tropical parasitoid wasps exposed by iterative integration of natural history. DNA barcoding. morphology and collections. *Proceedings of national academy of sciences, USA* 105: 12359-12364.
36. Borisenko AV, Lim BK, Ivanova NV, Hanner RH, Hebert PDN (2008) DNA barcoding in surveys of small mammal communities: a field study in Suriname. *Molecular Ecology Resources* 8: 471-479.
37. Kerr KCR, Lijtmaer DA, Barreira AS, Hebert PDN, Tubaro PL (2009) Probing evolutionary patterns in neotropical birds through DNA barcodes. *Plos One* 4: e4379.
38. Dewalt RE (2011) DNA barcoding: a taxonomic point of view. *Journal of the North American Benthological Society* 30: 174-181.
39. Szpila K (2012) Key for identification of European and Mediterranean blowflies (Diptera: Calliphoridae) of medical and veterinary importance – adult flies. Gennard D (ed.) *Forensic entomology, an introduction* 2: 77-81.
40. Nazni WA, Jeffery J, Heo CC, Chew WK, Lee HL (2011) Illustrated keys to adult flies of forensic importance in Malaysia. *Institute for Medical Research, Kuala Lumpur*.
41. Wallman JF (2001) A key to the adults of species of blowflies in southern Australia known or suspected to breed in carrion. *Medicine Veterinary and Entomology* 15: 433-437.
42. Kurahashi H, Benjaphong N, Omar B (1997) Blow flies (Insecta: Diptera: Calliphoridae) of Malaysia and Singapore. Singapore: School of Biological Sciences. University of Singapore.
43. Sukontason KLP, Narongchai P, Sripakdee D, Bhonchu N, Chaiwong T, et al. (2005) First Report of Human Myiasis caused by *Chrysomya megacephala* and *Chrysomya rufifacies* (Diptera: Calliphoridae) in Thailand, and Its Implication in Forensic Entomology. *Journal of Medicine and Entomology* 42: 702-704.
44. Turchetto M, Vanin S (2004) Forensic evaluations on a crime case with monospecific necrophagous fly population infected by two parasitoid species. *Aggrawal's Internet Journal of Forensic Medicine and Toxicology* 5: 12-18.
45. Carvalho LML, Linhares AX (2001) Seasonality of insect succession and pig carcass decomposition in a natural forest area in Southeastern Brazil. *Journal of Forensic Sciences* 46: 604-608.
46. CJB Carvalho, Ribeiro PB (2000) Key to the identification of species of Calliphoridae (Diptera) of southern Brazil. *Brazilian Journal of Veterinary Parasitology* 9: 169-173.
47. Rosa TA, Babata MLY, Sousa CM, Sousa DD, Mello-Patiu CA, et al. (2009) Diptera of forensic interest in two Cerrado vegetation profiles in Uberlândia, MG. *Neotropical Entomology* 38: 859-866.
48. Souza ASB, Kirst FD, Krüger RF (2008) Insects of forensic importance from Rio Grande do Sul State in southern Brazil. *Revista Brasileira de Entomologia* 52: 641-646.
49. Costa O, Mello-Patiu J, Lopes CA, Maria S (2001) Diptera associated with human corpses in the place of death in the state of Rio de Janeiro, Brazil. *Bulletin of the National Museum. New Series Zoology* 464: 1-6.
50. Carvalho LML, Thyssen PJ, Linhares AX, Goof ML (2004) Observations on the Successional Patterns of Necrophagous Insects on a Pig Carcass in an urban area of Southeastern Brazil. *Anil Aggrawal's Internet Journal of Forensic Medicine and Toxicology* 5: 33-39.
51. Chin HC, Marwi MA, Jeffery J, Omar B (2008) Insect succession on a decomposing piglet carcass placed in a man-made freshwater pond in Malaysia. *Tropical Biomedicine* 25: 23-29.
52. Salem AM, Adham FK, Picard CJ (2015) Survey of the Genetic Diversity of Forensically Important *Chrysomya* (Diptera: Calliphoridae) from Egypt. *Journal of Medical Entomology* 52: 320-328.
53. Gomes L, Von Zuben CJ, Govoni E (2004) Behavior of radial dispersion post larval food in blowflies *Chrysomya* genus (Diptera: Calliphoridae): search for new sources of food. *Entomologia Vector & Vectores* 9: 115-132.
54. Barros RM, Mello-Patiu CA, Pujol-Luz JR (2008) Sarcophagidae (Insecta, Diptera) associated with the decomposing carcasses of *Sus scrofa* Linnaeus (Suidae) in cerrado area of Federal District, Brazil *Revista Brasileira de Entomologia* 52: 606-609.
55. Barbosa TM, Mello-Patiu CA, Vasconcelos SD (2015) Flesh fly (Diptera: Sarcophagidae) survey on coastal environments in northeastern Brazil: new records and notes on the expanded geographical distribution. *Entomotropica* 20: 112-117.
56. Akbarzadeh K, Wallman JF, Sulakova H, Szpila K (2015) Species identification of Middle Eastern blowflies (Diptera: Calliphoridae) of forensic importance. *Parasitology Researcher* 114: 1463-1472.
57. Oliveira-da-Silva A, Ale-rocha R, Rafael JA (2006) Bionomics of the immature stages of two species of *Peckia* (Diptera, Sarcophagidae) in pig decaying in a forest area in northern Brazil. *Journal of Entomology* 50: 524-527.
58. ITIS Report (2011) *Phaenicia*. Integrated Taxonomic Information System.
59. Kosmann C, Mello RP, Harteirreiten-Souza ES, Pujol-Luz JR (2013) A List of Current Valid Blow Fly Names (Diptera: Calliphoridae) in the Americas South of Mexico with Key to the Brazilian Species. *EntomoBrasilis* 6: 74-85.
60. Rognes K (1991) Blowflies (Diptera, Calliphoridae) of Fennoscandia and Denmark. *Fauna Entomologica Scandinavica* 24: 90.